Central Asia Regional Trialogue on Land Degradation, Biodiversity and Climate Change

9-11 October 2019 | Almaty, Kazakhstan





How great is the problem? Status and Trends for Pollinators and Food Security

Hien T. Ngo (IPBES Secretariat)









Pollination Assessment Report & Process

- 2 year process (2014-2016)
- First output of IPBES
- 2 Co-chairs (Drs Simon Potts & Vera Imperatriz-Fonseca)
- 19 Coordinating Lead Authors
- 41 Lead Authors
- 14 Review Editors
- 35 Contributing Authors
- TOTAL: 76 experts










Output

Scientific literature review and Indigenous and local knowledge, 550pgs













Multiple Value of Pollinators

- Pollination of world's wild flowering plant species avg. almost 90% depend on pollinators (94% in tropical communities)
- Global food crops (production, yield and quality) (3/4 of the world's leading food types and 30% by volume) + regional crop economies rely (to varying degrees) on animal pollinators = <\$577 bn (USD/yr)!
- Healthy human diets and nutrition depend on pollinator dependent crops
- Livelihoods beekeeping and honey hunting & pollination services and food production
- medicines, biofuels (e.g. canola and palm oil), fibres (e.g., cotton and linen) construction materials (timbers), musical instruments, arts and crafts, recreational activities and as sources of inspiration for art, music, literature, religion, traditions, technology and education
- **globally significant heritage**, as symbols of identity, as aesthetically significant landscapes and animals, in social relations, for education and recreation and in governance interactions.
- Alhagi honey medicinal honey Turkmenistan



Man of Bicorp Valencia, Spain, 6000 BC



Multiple Value of Pollinators –Central Asia


- 14% of total agricultural crop output in Central Asia is dependent on pollination services (background document)
- Apples, strawberries, cucumbers, and tomatoes, pears, apricots, melons, watermelons etc.





The importance of pollinators

Percentage of production loss due to pollinator loss in leading global crops





















Pollinator-Mediated Crops & Nutrients in the Human Food Supply

- Staple crop production (e.g. cassava, corn, potato, rice, wheat, yam) = wind-pollinated, self-pollinated, or vegetatively propagated crops, they provide the majority of calories in the human diet = poor sources of micronutrients.
- Crop plants that depend fully or partially on animal pollinators contain >90% of vitamin
 the whole quantity of Lycopene and almost the full quantity of the antioxidants b-cryptoxanthin and b-tocopherol, the majority of the lipid, vitamin A (>70%) and related carotenoids (98%), calcium and fluoride, and a large portion of folic acid (55%)
- **58% of calcium** and **62% of fluoride** are derived from plants with marginally yield increase due to animal pollination, such as beans, but also strongly pollinator dependent plants such as fruits and nuts (including almonds) these minerals are crucial for development of teeth and bones and prevention of osteoporotic fracture risk
- 29% of iron is derived from pollinator-dependent crops, with 6% yield increase due to animal pollination – plant sources of iron crucial to human health
 - •Animal-pollinated crops contain the majority of the available dietary lipid, vitamin A, C and E, and a large portion of the minerals calcium, fluoride, and iron worldwide
 - •Ongoing pollinator decline may thus exacerbate current difficulties of providing a nutritionally adequate diet for the global human population.



Fractional dependency of micronutrient production on pollination





Pollinator-dependent crops, nutrition/health and Central Asia

- Vitamin A is one of the most prevalent deficiencies worldwide and is responsible for up to 500,000 annual cases of irreversible blindness in children worldwide
- More than 70% of vitamin A and 98% of each of the carotenoids cryptoxanthin (provitamin A) and lycopene are found in crops that are animal-pollinated
- As much as 50% of the production of plant-derived sources of vitamin A requires pollination throughout much of Southeast Asia, whereas other essential micronutrients such as iron and folate have lower dependencies, scattered throughout Africa, Asia and Central America.
- Micronutrient deficiencies are three times as likely to occur in areas of highest pollination dependence for vitamin A and iron, suggesting that disruptions in pollination could have serious implications for the accessibility of micronutrients for public health
- These regions of high nutritional vulnerability are understudied in the pollination literature, and should be priority areas for research related to ecosystem services and human well-being.



The diversity of pollinators

- flies, butterflies, moths, wasps, beetles, thrips, birds, bats and other vertebrates.
- BEES! 20,100 species of bees; most are wild




Apis mellifera pomonella Endemic honey bees of the Tien Shan Mountains in Central Asia





Global Trends

- Wild pollinators have declined in occurrence and diversity (and abundance for certain species) at local and regional scales in North West Europe and North America.
- Although a lack of wild pollinator data (species identity, distribution and abundance) for Latin America, Africa, Asia and Oceania preclude any general statement on their regional status, local declines have been recorded.
- Wild insects are extremely important pollinators and can pollinate many crops more effectively than managed bees
- The IUCN Red List assessments indicate that 16.5 per cent of vertebrate pollinators are threatened with global extinction (increasing to 30 per cent for island species). **There are no global Red List assessments specifically for insect pollinators.
- Regional and national assessments indicate high levels of threat for some bees and butterflies.





Trends in managed pollinators

 Global: The number of managed western honey bee hives has increased over the last five decades, even though declines have been recorded in some European countries and North America

•HONEYBEE TRENDS (1960s-2010)

Globally managed hives increased ~ 50%

The volume of production of pollinator dependent crops increased by 300%

Global agricultural is now twice as pollinatordependent



DRIVERS OF POLLINATOR AND POLLINATION CHANGE

- The abundance, diversity and health of pollinators and the provision of pollination are threatened by direct drivers that generate risks to societies and ecosystems.
 - •Threats include:
- o land-use change
- o intensive agricultural management and pesticide use
- o environmental pollution
- o invasive alien species
- o Pathogens
- Climate change




PESTICIDES

- Sulfoximine pesticides
- Reduced impacts in non-agricultural settings
 - Nanoparticle pesticides
 - Increasing fungicide use

CLIMATE CHANGE

- Extreme weather events
- Altered pathogen epidemiology

HABITAT LOSS AND HOMOGENISATION



Destruction of bat roosts

PARASITES AND PATHOGENS

- New RNA viruses
- Reduced pollinator richness drives epidemics
- Pollinators as disease vectors

NOVEL INDIRECT DRIVERS

- O P Increased diversity of managed pollinators
- Cutting pollinators out of food production
- Impacts of IPBES pollinators assessment



INVASIVE SPECIES

Invasive bees in Asia



 The seasonal loss of western honey bees in Europe and North America varies strongly by country, state and province and by year, but in recent decades (at least since the widespread introduction of Varroa) has often been higher than the 10-15 per cent that was previously regarded as normal.

Data for other regions of the world is largely lacking

- movement of *Apis* hives results in a spill over of pathogens both to this species, in the case of *Varroa* and from this species to wild pollinators like deformed wing virus
- Varroa mites, a key parasite of honey bees, have developed resistance to some chemical treatments so new treatment options are required



Beekeeping revitalization project takes off in Azerbaijan – case study in Azerbaijan's Caucasus Mountains & FAO

- Azeri beekeepers populated their apiaries with native Caucasian honeybees, which are well-known a strong and industrious honey producer.
- But after a deadly parasite devastated state-managed apiaries (early 1980s) colonies were restored with a different subspecies of bees that came from the country's southern region
- the new bees had a superior reproductive ability, they were weak honey producers compared to the Caucasian bee.
- Over time, the two subspecies hybridized = mixed-race bee that produced less honey per hive than before
- Azerbaijan requested FAO assistance through the Turkish Partnership Programme Trust Fund, which has dedicated US\$ 200,000 to the project and made Turkish apiculture specialists available to train Azeri beekeepers.
- By increasing Azerbaijan's honey production, FAO experts expect that more honey will be available at lower prices in city markets.
- Back in the Caucasus Mountains, beekeepers will see increased rural employment and income—both important steps toward reducing rural poverty.



Land Use Change

- Habitat destruction
- fragmentation and degradation
- conventional intensive land management practices:
- high use of agrochemicals
- intensively performed tillage, grazing or mowing
- By 2030, the area of agricultural land is expected to increase a further 10%, mainly in the developing world





Central Asia Regional Trialogue on Land Degradation, Biodiversity and Climate Change




FERTILISER, PESTICIDE USE

- Risk to pollinator depends on the combination of toxicity and the level of exposure, which varies geographically with the compounds used and the scale of land management and habitat in the landscape.
- Pesticides (sp. Insecticides) have a broad range of lethal and sublethal effects on pollinators under controlled experimental conditions.
- It is currently unresolved how sublethal effects of pesticide exposure recorded for individual insects affect colonies and populations of managed bees and wild pollinators, especially over the longer term.
- There is evidence from a recent study that shows impacts of neonicotinoids on wild pollinator survival and reproduction at actual field exposure. Evidence, from this and other studies, of effects on managed honey bee colonies is conflicting

For many other countries (e.g., in Africa and Asia) data are incomplete or absent.



FERTILISER, PESTICIDE USE





Genetically Modified Organisms (GMOs)

- Are herbicide tolerance (HT) or insect resistance (IR) (traits); reduced weed populations accompany HT crops which decrease food resources for pollinators. Insect resistant (IR) crops can result in the reduction of insecticide use - the reduction in insecticide use could reduce pressure on non-target insects.
- Risk assessments required for the approval of genetically modified organism (GMO) crops in most countries do not adequately address the direct sublethal effects due to lack of data

Invasives

 can disrupt native pollinator communities by outcompeting indigenous insects for resources or by spreading pests and disease - there is little available evidence that alien plants are detrimental to pollinator diversity

Diseases and pests

- Bees suffer from a broad range of parasites, including *Varroa* mites in western and eastern honey bees.
- Emerging and re-emerging diseases are a significant threat to the health of bees, especially when they are managed commercially.



Climate change

- Climate change is anticipated to bring about changes in rainfall distribution, wind patterns, temperature, air pollution and occurrence of extreme weather events, among other environmental changes
- The effects of climate change on plant-pollinator interactions are still mostly unknown and the indirect effects upon interacting species and networks of species are poorly represented in the literature
- Plant and pollinator ranges are shifting/moved their ranges, altered their abundances and shifted their seasonal activities in response to observed climate change over recent decades <u>A recent analysis: bumble bees appear to be undergoing range contractions as climate changes across Europe and North America</u>
- Differential migration rates of co-occurring plants and insects as a result of changing climatic conditions may lead to a spatial dislocation of processes like pollination.
- climate change may alter the synchrony between plant flowering and pollinator flight periods = PHENOLOGICAL MISMATCHES = probably contribute to pollinator losses that subsequently disrupt pollination of plants



Climate change scenarios

- Under all climate change scenarios for the second half of the 21st century
 - •(i) pollinator community composition is expected to change as a result of decreases in the abundance of some species and increases in others
 - •(ii) the seasonal activity of many species is predicted to change differentially, potentially disrupting life cycles and species interactions between plants and pollinators
 - •Impacts of ongoing climate change on pollinators and pollination services to agriculture may not be fully apparent for several decades, owing to a delayed response in ecological systems.
 - •Adaptive responses to climate change include increasing crop diversity and regional farm diversity and targeted habitat conservation, management or restoration.
 - •The effectiveness of adaptation efforts at securing pollination under climate change is untested



International actions as a result of pollination assessment Convention on Biological Diversity (CBD)

The Thirteenth meeting of the Conference of the Parties (COP 13) (Cancun, Mexico, December 2016) adopted decision XIII/15 on the Implications of the IPBES assessment on pollinators, pollination and food production for the work of the Convention, in which the Conference of the Parties welcomed the IPBES summary for policymakers of the thematic assessment on pollinators, pollination and food production and endorsed its key messages.







International actions as a result of pollination assessment Convention on Biological Diversity (CBD)

- PROMOTING POLLINATOR-FRIENDLY HABITATS
- IMPROVING THE MANAGEMENT OF POLLINATORS, AND REDUCING RISK FROM PESTS, PATHOGENS AND INVASIVE SPECIES
- REDUCING RISK FROM PESTICIDES, INCLUDING INSECTICIDES, HERBICIDES AND FUNGICIDES
- ENABLING POLICIES AND ACTIVITIES
- RESEARCH, MONITORING AND ASSESSMENT





Declaration - Coalition of the Willing on Pollinators

 The Netherlands announced at the high level segment of COP-13, the launch of a new initiative, called the Coalition of the Willing on

Pollinators.

Signatories include:

• France, Germany,

- the United Kingdom,
- Belgium, Austria,
- Denmark,
- Luxembourg, Finland,
- Spain, the Netherlands,
- Uruguay, Peru and
- Slovenia (n=13)
- *Ethiopia signed the Declaration on the Coalition of the Willing on Pollinators (2017)
- Total membership = Coalition has 24 member countries







Many National Initiatives Established or in development

England
Scotland
Wales
All Ireland
France
Switzerland

Netherlands

Norway

USA

Canada

Brazil

Argentina

Colombia

South Africa

India

Republic of Korea

EU and global and others





 Awareness and communication opportunities: World Bee Day https://www.worldbeeday.org/en/bee-with-us.html



Capacity Building & mainstreaming:

BES-Net Trialogue (Sarajevo, Bosnia and Herzegovina) 18-20 October 2017



BES-Net Trialogue (Santo Domingo, Dominican Republic) Anglophone Africa (Nairobi, Kenya)

More Initiatives

- National pollinator strategies (to be developed)
- EU Pollinators Initiative
- The International Pollinator Initiative Plan of action 2018-2030 based on IPBES

Asian Pollinator Initiative**

CBD – Decision 14/6 - Adopts the Plan of Action 2018-2030 for the International Initiative for the Conservation and Sustainable Use of Pollinators as contained in annex I to the present decision, for implementation according to national legislation and national circumstances



Asian Pollinator Initiative

27 - 29th February, 2020 at University of Calcutta, Kolkata, India.

The meeting is being organized jointly by the Centre for Pollination Studies (CPS), University of Calcutta, National Centre for Biological Sciences (NCBS), Bengaluru and University of Agricultural Sciences (UAS), Bengaluru in collaboration with the Zoological Survey of India.





Conclusions

- Pollinator loss is an urgent problem
- Food Security: High policy relevance (impact on food production)
- Nutritional security: pollinators improve yields for crops that contribute nutrients to the food supply
- Economic, Ecological, social and cultural importance
- Impact is still limited by the lack of global policy uptake and implementation effort
- Globally we share many of the broad challenges and these will need locally and regionally developed solutions to enhance pollinators and pollination services
- Regional and national initiatives have a critical role to play in sharing knowledge, building capacity and supporting the development of better policies and practices





IPBES Secretariat, UN Campus Platz der Vereinten Nationen 1, D-53113 Bonn, Germany secretariat@ipbes.net





How to become a member of IPBES?

- intention should kindly be communicated to the IPBES secretariat by way of a formal letter
- (2) issued by or on behalf of the authority that would issue the credentials for your national delegation to a session of the IPBES Plenary, in other words by your Head of State or Government or minister for foreign affairs, consistent with your national policies and law
- (3) letter should also please identify, and provide official contact details for your Government's National Focal Point for IPBES
- (4) all contributions to the IPBES trust fund are entirely voluntary

Email: Satomi Yoshino - satomi.yoshino@ipbes.net